

XP 002152630

AN - 1997-234560 [21]

AP - SU19925042406 19920515

CPY - PUKH-I

DC - B04 D16 S03

FS - CPI;EPI

IC - G01N33/52 ; G01N33/557

IN - PUKHOV K I; PUKHOVA YA I; ZVEGINTSEV S A

MC - B04-F04 B11-C07B4 B12-K04 D05-H09

- S03-E14H S03-E14H4

M1 - [01] M423 M750 M903 N102 Q233 V600 V612 V754

M6 - [02] M903 P831 Q233 R160 R514 R611 R625 R635

PA - (PUKH-I) PUKHOVA YA I

PN - RU2067766 C1 19961010 DW199721 G01N33/52 005pp

PR - SU19925042406 19920515

XA - C1997-075171

XIC - G01N-033/52 ; G01N-033/557

XP - N1997-193985

- AB - RU2067766 Evaluation of the functional activity of immunocompetent cells (ICC) in the whole blood system involves selection of blood samples, recording the level of chemiluminescence in the presence of an inductor and a chemiluminescence probe, with subsequent calculation of an evaluation criterion E. The chemiluminescence light sum S, equal to the area under the chemiluminescence reaction curve (CLR), and the time during which 50% of the CLR light sum is realised (Tc), are recorded. The evaluation criterion E characterizing the degree of difference between the two CLR, i.e. that for the patient and that for the norm, is the difference between these two curves which can be represented on a plane in the form of 2 points in the S, Tc coordinate system. This is calculated as the normalised square of the distance between these two points to the square of the distance from the origin of coordinates (Tv, S) to the point CLR1 (point of the norm) from the formula $E = 100 (Tc(pat) - T(nor))^2 + (S(pat) - S(nor))^2 / (Tc(nor)^2 + S(nor)^2)$, where the parameters of the mean statistical CLR curve are taken as the norm: for practically healthy patients. Additional kinetic CLR characteristics are then calculated in the S, Tc coordinate system as the difference (S (pat)-S(nor)) and (Tc(pat)-Tc(nor)), from which when S pat is greater than S nor + sigma, a judgement is made that the CLR activity of the patient is higher than the upper limit of the norm, and if S pat is less than S nor - sigma, a judgement is made that the CLR activity is below the lower limit of the norm. If Tc pat is less than Tc nor - sigma, a 'left shift' judgement is made, indicating an increase in the CLR above the upper limit of the norm, and if Tc pat is greater than Tc nor + sigma, a 'right shift', indicating a decrease in the CLR rate below the lower limit of the norm.
- USE - The method is useful in clinical medicine, experimental and clinical immunology and general pathology, for early diagnosis of immunodeficient states, for monitoring the functional activity of immunocompetent cells in immuno-correction, in testing pharmacological preps. and in evaluating the effectiveness of intensive therapy.
 - ADVANTAGE - The method is simpler, more accurate and more informative

than previous methods.

- (Dwg.0/4)

IW - EVALUATE FUNCTION ACTIVE IMMUNO COMPETENT CELL WHOLE BLOOD SYSTEM

RECORD LIGHT SUM CHEMILUMINESCENT TIME PER CENT LIGHT SUM REALISE

IKW - EVALUATE FUNCTION ACTIVE IMMUNO COMPETENT CELL WHOLE BLOOD SYSTEM

RECORD LIGHT SUM CHEMILUMINESCENT TIME PER CENT LIGHT SUM REALISE

INW - PUKHOV K I; PUKHOVA YA I; ZVEGINTSEV S A

NC - 001

OPD - 1992-05-15

ORD - 1996-10-10

PAW - (PUKH-I) PUKHOVA YA I

TI - Evaluating functional activity of immuno-competent cells in whole
blood system - with recording of the light sum of chemiluminescence
and time during which 50 per cent of the light sum is realised